

The chiral symplectic universality class

Yoichi ASADA*, Keith SLEVIN and Tomi OHTSUKI¹

Department of Physics, Graduate School of Science, Osaka University, 1-1 Machikaneyama, Toyonaka, Osaka 560-0043, Japan

¹*Department of Physics, Sophia University, Kioi-cho 7-1, Chiyoda-ku, Tokyo 102-8554, Japan*

(Received February 2, 2008)

KEYWORDS: Anderson localization, chiral symmetry, spin-orbit coupling

1. Chiral symmetry

About a decade ago Gade¹⁾ investigated the effect of chiral symmetry in a nonlinear sigma model (Gade model). She showed that this symmetry causes the localisation length and the density of state to diverge at the band center in 2D. This behaviour at the band center is different from that found in the “standard” universality classes of Anderson localisation.

If a Hamiltonian H anti-commutes with a unitary matrix U then H has chiral symmetry. In this case energy eigenvalues occur in pairs of opposite sign with pairs of eigenstates ψ_n and $U\psi_n$. When there is no diagonal disorder the Hamiltonian may have chiral symmetry. This occurs if we can divide the system into two sublattices such that nearest neighbour sites belong to different sublattices. If this is possible, the transformation U which changes the sign of the wave function only on one sublattice anti-commutes with the Hamiltonian. If the total number of lattice sites is odd, chiral symmetry guarantees the existence of zero energy eigenstate(s), or zero mode(s). Such states, if they exist, have special properties e.g. they are supported only on one sublattice. The precise number of zero modes depends on the difference of the number of sites in each sublattice.²⁾

One example of a system with chiral symmetry is the random magnetic flux model. Anomalous behaviour is observed at the band center.³⁾ The results indicate that the entire band is localised except for a critical state at the band center.

2. The SU(2) model

Here we report a numerical investigation of localisation in the SU(2) model⁴⁾ without diagonal disorder. The Hamiltonian of the SU(2) model has time reversal symmetry but spin rotation symmetry is broken by a random spin-orbit coupling. Thus we expect that states will belong either to the standard symplectic class or the chiral symplectic class. This model is different from the random magnetic flux model in that the zero mode is surrounded by a metallic phase, not by an insulating phase.

The Hamiltonian of the SU(2) model describes non-interacting electrons on a simple square lattice with near-

est neighbour SU(2) random hopping

$$H = - \sum_{\langle i,j \rangle, \sigma, \sigma'} U(i, j)_{\sigma\sigma'} c_{i\sigma}^\dagger c_{j\sigma'} \quad (1)$$

where $c_{i\sigma}^\dagger$ ($c_{i\sigma}$) denotes the creation (annihilation) operator of an electron at site i with spin σ . We distribute hopping matrices randomly and independently with uniform probability on the group SU(2):

$$U(i, j) = \begin{pmatrix} e^{i\alpha_{ij}} \cos \theta_{ij} & e^{i\gamma_{ij}} \sin \theta_{ij} \\ -e^{-i\gamma_{ij}} \sin \theta_{ij} & e^{-i\alpha_{ij}} \cos \theta_{ij} \end{pmatrix} \quad (2)$$

where α and γ are uniformly distributed in the range $[0, 2\pi)$, and θ distributed in the range $[0, \pi/2]$ according to the probability density, $P(\theta)d\theta = \sin(2\theta)d\theta$.

The SU(2) model has a mobility edge at $E_c \simeq 3.253$.⁴⁾ In what follows we focus attention on the properties of the model near the band center at $E = 0$. We calculate the localisation length on a quasi-1D strip and attempt to extrapolate to the 2D limit.

3. Behaviour of the localisation length

We consider a quasi-1D strip whose width is L and calculate the localization length λ at arbitrary energy E with the transfer matrix method.⁵⁾ In the transverse direction we impose either fixed boundary conditions (FBC) or periodic boundary conditions (PBC). The Hamiltonian has chiral symmetry except when PBC are imposed on a system whose width is odd. In this latter case chiral symmetry is broken. We analyse the dependence of the re-normalized localization length $\Lambda = \lambda/L$ on L .

The re-normalized localization length Λ at $E = 0.0$ and $E = 0.5$ as a function of L are shown in Figures 1 and 2 respectively. The width L ranges from 7 to 128. We performed $10^6 \sim 10^8$ transfer matrix multiplications to obtain data with an accuracy of 0.05% to 2%.

Figure 1 shows the results at the band center $E = 0.0$. (Note that for odd L and FBC the localisation length λ diverges⁶⁾ so no data is presented for this case.) A striking dependence on the parity of L is observed. As a function of L , Λ approaches a strongly parity dependent constant value for $L \rightarrow \infty$. For even L this asymptotic value is boundary condition independent. To analyse the data in detail, we fit the data to

$$\Lambda = a + cL^{-y}. \quad (3)$$

* asada@presto.phys.sci.osaka-u.ac.jp




Fig. 1. Λ vs L at $E = 0.0$. The solid lines are the fit.

The results are tabulated in Table I. For even L the estimates of the asymptotic value a are the same within numerical accuracy. For odd L and PBC the asymptotic value is much larger than for even L . The L independent behaviour of Λ at large L is typical of a critical point and may indicate that band center is critical in this system.




Fig. 2. Λ vs L at $E = 0.5$. The solid lines are the fit.

In contrast at $E = 0.5$ (Figure 2) the parity dependence is negligible (except at small L under PBC). Though a much strong boundary condition dependence is now observed, it seems this will disappear in the limit of large L . For PBC the data can be fitted by

$$\Lambda = a + b \ln L. \quad (4)$$

and for FBC by

$$\Lambda = a + b \ln L + cL^{-y}. \quad (5)$$

The results are tabulated in Table II. In the limit of sufficiently large L it seems that a boundary condition independent logarithmic increase of Λ will be recovered. Note in particular, by reference to Table II, that the

estimated (asymptotic) slopes for both curves in Figure 2 are in agreement.

To compare this behaviour with that in the metallic phase of the standard symplectic class, we also performed simulations of the $SU(2)$ model with diagonal disorder by adding a term

$$H_d = \sum_{i,\sigma} \epsilon_i c_{i\sigma}^\dagger c_{i\sigma} \quad (6)$$

to the Hamiltonian (1). Here ϵ_i distributed randomly

Table I. The best fit estimates of fitting parameters at $E = 0.0$ with their 95% confidence intervals. Q is the goodness of fit probability.

	Q	a	c	y
FBC,Even	0.8	$1.39 \pm .01$	$2.6 \pm .4$	$1.12 \pm .08$
PBC,Even	0.8	$1.381 \pm .002$	$-2 (-6, -0.8)$	$2.2 \pm .5$
PBC,Odd	0.8	$4.48 \pm .03$	3.2 ± 1.3	$1.2 \pm .2$

Table II. The best fit estimates of fitting parameters at $E = 0.5$ with their 95% confidence intervals. Q is the goodness of fit probability. In fitting we eliminated the data for $L \leq 8$.

	Q	a	b	c	y
PBC	0.1	$4.14 \pm .04$	$0.653 \pm .013$		
FBC	0.2	5.0 ± 1.6	$0.6 \pm .3$	12 ± 6	$1.0 \pm .3$

and independently with uniform probability on the interval $[-1, 1]$. The diagonal disorder breaks chiral symmetry and the system belongs to the standard symplectic class. The re-normalized localization length Λ is again a logarithmic function of L and we are able to fit the result by (4). We also get an estimate of the coefficient $b = 0.64 \pm .02$ in the $SU(2)$ model which is consistent with those in Table II. This indicates that the states away from the band center in the $SU(2)$ model without diagonal disorder are metallic and belong to the standard symplectic universality class.

- 1) R. Gade, Nucl. Phys. B **398** 499 (1993).
- 2) P. W. Brouwer, E. Racine, A. Furusaki, Y. Hatsugai, Y. Morita, and C. Mudry, Phys. Rev. B **66** 014204 (2002).
- 3) A. Furusaki, Phys. Rev. Lett. **82** 604 (1999).
- 4) Y. Asada, K. Slevin and T. Ohtsuki, Phys. Rev. Lett. **89** 256601 (2002).
- 5) A. MacKinnon and B. Kramer, Z. Phys. B **53**, 1 (1983).
- 6) P. W. Brouwer, C. Mudry, B. D. Simons, and A. Altland, Phys. Rev. Lett. **81** 862 (1998).